arXiv:1304.2710v1 [cond-mat.soft] 9 Apr 2013

Helixing Transition in Filaments under Torque

Cristiano Nisoli and A. R. Bishop
Theoretical Division and Center for NonLinear Studies, Los Alamos National Laboratory, Los Alamos NM 87545 USA

(Dated: April 10, 2013)

We show that a filament in thermal equilibrium subjected to torque τ and a large enough tension undergoes a phase transition of infinite order when the torque is twice the temperature T: for very large values of the ratio τ/T, the polymer's average supercoiling responds linearly to the applied torque, but it tends to zero exponentially fast as the ratio reaches the critical value $\tau/T = 1/2$. When approaching the transition from below, correlation lengths tend exponentially to infinity, which suggests an essentially topological nature. At room temperature, this transition happens at a torque ~ 2 pN×nm, and it should be detectable in carefully performed single molecule manipulation experiments.

PACS numbers: 82.35.Pq,05.70.Fh, 87.14.E-, 87.14.G-

The thermodynamics of filaments has been the center of extensive study over the last four decades, since the pioneering works of Edwards, De Gennes and others [1, 2] started what is now called soft condensed matter physics. Motivated by practical reasons and advances in polymer science, by e.g. Kuhn and Flory [3], research in polymer physics initially concentrated on the study of collective statistical behavior: solution, gelation, networks, entangling and unentangling, rheology, *etc.* [2].

Biological and technological interest in highly structured polymers, most notably DNA, have motivated experimental research into their behavior as single molecules under precisely controlled mechanical manipulation. Direct single molecule manipulation of DNA, proteins, and other more or less complex polymers [4] has opened the field to the analysis of entirely new phenomena, such as writhing and load/temperature induced denaturation. Meanwhile, advances in nanotechnology have shown how single polymers and DNA can be compose into novel materials [5], whereas novel inorganic nano-filaments such as carbon nanotubes were also introduced [6]. The confluence of theoretical, experimental, and technological developments has made it timely to analyze the behavior of single (rather than many) polymers under load; much effort [7–11] has concentrated on understanding single molecule experiments for DNA in particular [12–14].

Here we show that an infinitely smooth helixing transition is expected in a long filament under applied load in a thermal bath. The transition is controlled by the ratio between applied torque τ and temperature T (measured as energy): for $T \ll \tau$, the polymer responds linearly to torque as it assumes an helical structure. For $T > \tau$ the order parameter (helicity) disappears and the structural response of the polymer to torque is cancelled by thermal fluctuations, as its generalized helical rigidity becomes infinite. The transition is of infinite order and characterized by an exponential divergence of correlation lengths.

Consider modeling the following experiment: a mesoscopic one-dimensional system, which for simplicity we take as a polymer with monomers of length a, is held under tension χ and subjected to a torque τ by magnetic or optical tweezers [4]. Held straight, under applied torque it will respond with an elastic change of the internal torsional angle. On the other hand, if the torque is large enough, one expects a structural change, in which the polymer assumes an helical configuration.

We consider a continuum energy model for this system. Suppose the tension is large enough that χa^2 exceeds the flexural rigidity [21]: then the dominant contribution to the energy will come from the change $\chi \Delta h$ in distance h between the ends by which the filament is held under tension χ. If, in a continuum limit, $\psi(s) = r(s) \exp[i\theta(s)]$ represents the deviation from the straight filament configuration in the plane perpendicular to it , and s is the intrinsic coordinate (see Fig. 1), then the contribution (per unit length) to the energy density from tension, neglecting subdominant terms, is $-\chi dh/ds = \chi(\dot{r}^2 + r^2\dot{\theta}^2)/2$. Here dh is the experimentally measured change in distance h between beads. The contribution from the torque is $-\tau\vec{t}\cdot\vec{d}$, where $\vec{t}$ is the unit vector tangent to the curve and $\vec{d}$ the Darboux vector. For configurations that wind around the experimental axis, it can be approximated as $-\tau\dot{\theta}$.

One arrives at the following expression for the energy per unit length

$$\mathcal{H} = \frac{1}{2}\chi\left(\dot{\rho}^2 + \rho^2\omega^2\right) - g(\rho)\tau\omega, \tag{1}$$

where $\rho \in [0, \infty)$, $\omega = \dot{\theta} \in (-\infty, \infty)$ are functions of the parameter $s \in [0, L)$, with L large in the thermodynamic limit. The regularization at small ρ, $g(\rho)$, is $g = 1$ when $\rho > \rho_0$ (a cutoff at small ρ, ρ_0 is of the order of the lateral size of the polymer), and $g = 0$ otherwise. A less rigorous choice does not affect the following conclusions. The cutoff $g(\rho)$ avoids unphysical configurations in which $\rho = 0$ yet $\omega \neq 0$. More importantly, it gives a lower bound to the energy (1). Physically it describes the breakdown of the model when the radius of the helix becomes comparable to the radius of the polymer, and couplings between



FIG. 1: The polymer held at its ends by a tension χ along the direction h and subject to a torque τ assumes a helical configuration (top right): the two-dimensional vector $\psi(s)$ (red arrow) describes the deviation from the straight filament configuration in the plane perpendicular to h, while s is the intrinsic coordinate. For flexural modes of the helix (bottom right, blue arrow), which are suppressed by tension and do not couple with the torque, ψ describes the deviation from the flexural mode. Top left: as $\tau/T \to 1/2+$ the average helicity $\langle\omega\rangle l$ (red solid line) tends to zero exponentially fast, while both the generalized helical rigidity $\langle k\rangle \equiv 1/\langle k^{-1}\rangle$ (dashed black line) and the correlation length ξ (black solid line) approach infinity with exponential behavior. Bottom left: in an experiment of single molecule manipulation one often measures the torque after having fixed the supercoiling σ. The transition is then revealed by plotting the torque *vs.* $\sigma-\tau/\mu$ to suppress the inner torsional variable of rigidity μ.

the helical gain and the inner torsional degree of freedom become relevant.

One can check that, if ρ is kept fixed, the energy in (1) is minimized by an helix of gain angle per unit length

$$\omega = k(\rho)^{-1}\tau, \tag{2}$$

where $k(\rho)$ is the linear density of helical rigidity of the structure per unit length (and therefore k^{-1} is the helical compliance) and normalized to temperature. We have:

$$k(\rho)^{-1} = \frac{g(\rho)}{\chi\rho^2}. \tag{3}$$

The energy of the helix is $\mathcal{H}[\rho,\omega(\rho)] = V(\rho)$, with

$$V(\rho) = -\frac{\tau^2}{2\chi}\frac{g(\rho)}{\rho^2}. \tag{4}$$

If there is no constraint on the radius ρ, the lowest energy structure corresponds to $\rho=\rho_0$ and helicity $\omega_0 = k(\rho_0)^{-1}\tau$. More realistic choices for the cutoff $g(\rho)$ accommodate the statics of different polymers but have no effect on the transition.

Let us now investigate the effect of thermal fluctuations on this structure. The term $-\tau\omega$ disregards the energy of polymer configurations winding around a different axis than the experimental one: nonetheless, one can always decompose these configurations in a slowly varying flexural mode–suppressed by tension–plus a helical mode described by our hamiltonian. The flexural modes (Fig. 1) are suppressed by the tension and do not couple to the torque, therefore representing an inconsequential background for our torque-driven transition.

In the presence of thermal fluctuations, entropy reduction is the cost of structure. Helical fluctuations , although favored by energy, should be entropically disadvantageous compared to planar ones. One can anticipate that this competition between entropy and energy might drive a transition.

We can gain some heuristic insight by computing the free energy of a helix of fixed angular gain $\bar\omega$. We sum over fluctuations *in* ρ *only* while maintaining $\bar\omega$ fixed, to obtain the partition function of an helix of gain $\bar\omega$, or $Z_{\bar\omega} = e^{-\frac{\tau\bar\omega L}{T}}\int[d\rho]\ e^{-\frac{\chi}{2T}\int_0^L(\dot\rho^2+\rho^2\bar\omega^2)ds}$, where $[d\rho] = \prod_{i=1}^{L/a} d\rho(ai)$ in the limit of small a. This path integral can be computed exactly by turning it into the trace of the density operator for the harmonic Schrödinger problem in ρ, which then projects on its lowest eigenvalue $|\bar\omega|/2$ in the thermodynamic limit of large L. Thus one finds the free energy density $f_{\bar\omega} = L^{-1}\ln Z$ for an helix of *fixed* helicity $\bar\omega$. By subtracting the equipartition free energy density (which is $T\ln(\chi/T)/2a$, obtained by setting $\bar\omega = 0$) one finds the extra free energy contributed by a structure of defined helicity $\bar\omega$,

$$\Delta f_{\bar\omega} = -\tau\bar\omega + T|\bar\omega|/2, \tag{5}$$

which corresponds to an energy reduction $-\tau\bar\omega$, but also an entropy reduction $\Delta s = -|\bar\omega|/2$. Their interplay presages a transition: for values of $|\tau|/T < 1/2$ helical structures are suppressed, as any helicity $\bar\omega \neq 0$ gives a positive contribution to the free energy. However when $|\tau| > T/2$, or equivalently $T < T_c = 2|\tau|$, the entropy cost of helicity can be offset by energy gain and helical structures of the same orientation of τ lower the free energy [22]. As we will see below, this heuristic argument correctly predicts the critical temperature $T_c = 2|\tau|$.

To obtain the exact nature of the transition we return to the full partition function

$$Z \propto \int[\rho d\omega d\rho]e^{-\beta\int_0^L \mathcal{H}ds}. \tag{6}$$

Since $\mathcal{H}$ in (1) is quadratic in ω, the helicity can be integrated out. This removes the ρ factor from the functional measure in (6) and provides an equipartition factor, returning

$$Z \propto \left(\frac{2\pi T}{\chi}\right)^{\frac{L}{2a}} \int [d\rho] e^{-\beta \int_0^L \left[\frac{\chi}{2}\dot{\rho}^2 + V(\rho)\right] ds}. \tag{7}$$

From (6) we can define the order parameter for the helicity

$$\langle\omega\rangle = T L^{-1} \partial_\tau \ln Z = \langle k^{-1}\rangle \tau, \tag{8}$$

where $\langle k^{-1}\rangle$ is the generalized helical compliance. Equation (8) can be rewritten as $\langle k^{-1}\rangle = 2TL^{-1}\partial_{\tau^2} \ln Z$ and then from (7) and (4) one obtains

$$\langle k^{-1}\rangle = \langle \frac{g(\rho)}{\chi\rho^2}\rangle = \langle k(\rho)^{-1}\rangle : \tag{9}$$

the generalized helical compliance is simply the thermal average of the "static" helical compliance given in (3).

The helical compliance is non-zero when the radius is finite. From (8) and (9), when $\langle\omega\rangle \neq 0$, we can picture the system as fluctuating around a helix of gain angle $\langle\omega\rangle$, and of radius given by $\langle\rho^{-2}\rangle = \chi\langle k^{-1}\rangle = \langle\omega\rangle/\tau$: consistent with our heuristic findings, the radius is confined when helical thermal fluctuations dominate.

The partition function in (7) can be expressed through standard path integral manipulation in terms of a trace

$$Z \propto \left(\frac{T}{\chi}\right)^{\frac{L}{a}} \mathrm{Tr}\left(e^{-L\hat{H}}\right) \tag{10}$$

of a one dimensional Schrödinger operator

$$\hat{H} = -\frac{T}{2\chi}\left[\frac{d^2}{d\rho^2} + g(\rho)\frac{(\tau/T)^2}{\rho^2}\right]. \tag{11}$$

[The factor of proportionality in (10) depends on boundary condition and is non-extensive in L. In the absence of truncation g the operator in (11) is unbounded from below, just like the energy in (1).]

In the thermodynamic limit of large L, the trace in (10) projects on the lowest bound state of eigenvalue ϵ_0—if there is one. The pre-factor $(T/\chi)^{L/a}$ in (10) only contributes two equipartition terms corresponding to the two oscillating variables ω and ρ and is independent of τ. Thus one has $\langle\omega\rangle = -T\partial_\tau\epsilon_0 \neq 0$, giving rise to helical structures of confined radius, as described above, whenever there is a discrete spectrum.

The operator in (11) corresponds to the well known Schrödinger problem of an attractive truncated inverse square potential. Its discrete spectrum is known to disappear when $|\tau|/T < 1/2$, which confirms the existence of a transition at $T_c = 2|\tau|$. Indeed since the contribution from the continuum spectrum in (10) is non-extensive in L, for $T > T_c$ the partition function reduces to equipartition alone, or $Z = (T/\chi)^{(L/a)}$, which is independent of τ, and thus $\langle\omega\rangle = 0$ above the transition: the helical structure is lost.

When $T \to 2\tau^-$ the 1-D Schrödinger problem of a truncated inverse square potential on a half line has been studied by Gupta and Rajeev [15] through renormalization methods, later extended to general dimension by Camblong *et al.* [16]. Indeed the problem is a long standing one of quantum mechanics, and has relevance in molecular physics [17], string theory [18], and polymer physics (in different context than ours) [19].

Defining $\nu^2 = (\tau/T)^2 - 1/4$, the remaining bound eigenvalue can be approximated in the limit $\nu \to 0^+$ [15] or $\tau/T \to (1/2)^+$ as

$$\epsilon_0 = -l^{-1} e^{-\frac{2\pi}{\nu}}[1 + O(\nu)], \tag{12}$$

with $l = \rho_0^2 e^{2\gamma}\chi/T$, and γ the Euler-Mascheroni constant. From (12) it follows that the helical order parameter

$$\langle\omega\rangle \simeq \frac{2\pi}{l}\frac{\tau}{T\nu^3} e^{-\frac{2\pi}{\nu}}\left[1 + O(\nu^2)\right] \tag{13}$$

disappears at transition with infinite smoothness (all its derivatives tend to zero), Fig. (1). The transition is therefore of infinite order, as one might anticipate given its topological nature. Clearly, at transition the generalized rigidity $\tau/\langle\omega\rangle$ approaches infinity exponentially fast and therefore, from (9), so does $\langle\rho^2\rangle$.

Finally, it is interesting to reason in terms of helical pitch, $2\pi/\langle\omega\rangle$. In particular its ratio with the radius of the helix, from (9) is $\propto \langle\omega\rangle^{-1/2}$. Therefore the pitch of the helix in units of the radius of the helix approaches infinity exponentially fast at transition, i.e. the helix becomes degenerate to a plane infinitely fast. Above the transition, as we have seen, only planar fluctuations contribute to the partition function.

Our previous heuristic treatment lead us to suspect that fluctuating helical configurations provide a dominant contribution to the partition function below the transition. In contrast, only planar oscillations contribute above the transition. Eq. (13) shows that these helical structures disappear very smoothly.

From (10) one has for the free energy density $f = T\epsilon_0 + a^{-1}T\ln(\chi/T)$. Neglecting the last term, which corresponds to equipartition, one finds from $s = -\partial_T f$

$$\Delta s = -\frac{1}{2}\langle\omega\rangle\left[1 + 2\nu^2 + O(\nu^3)\right], \tag{14}$$

which shows that close to transition, the principal part of the entropy reduction is proportional to $-\langle\omega\rangle/2$. This is the same result we obtained from (5) for the structural cost in entropy for a helix of *uniform* gain, suggesting that at transition the order parameter tends to uniformity as it disappears: its fluctuations tend to zero, while their correlation length approaches infinity. The first statement is obtained immediately by differentiating the expression for $\langle\omega\rangle$ in (13) with respect to τ. We demonstrate the second one below.

The infinite order character of the transition as well as its topological nature should also be evidenced in an exponentially smooth disappearance of the correlation lengths. These can be computed as follows. We introduce a varying external field $\tau(s) = \tau + \eta(s)$ with τ uniform, as before. Then we compute the functional $Z[\eta(s)]$ by summing over all the trajectories in $\omega(s)$, $\rho(s)$ as in (6). Clearly the helical correlation function $G(s_1, s_2) = \langle\omega(s_1)\omega(s_2)\rangle - \langle\omega\rangle^2$ is given by [20]

$$G(s_1, s_2) = T^2 \frac{\delta^2 \ln Z[\eta]}{\delta\eta(s_1)\delta\eta(s_2)}|_{\eta=0}. \tag{15}$$

The new partition function $Z[\eta]$ for $\eta(s) \neq 0$ is still given by a trace as in (10), but with the replacement $L\hat{H} \to L\hat{H} + \hat{\phi}\int_0^L \eta(s)ds$, with $\phi(\rho) = g\tau/(T\chi\rho^2)$. Using imaginary time $s-$dependent perturbation formalism, one obtains through standard calculations

$$G(s_1, s_2) = \frac{T^2}{2}\int_0^\infty dk |\langle\epsilon_0|\hat{\phi}|k\rangle|^2 e^{-\Delta\epsilon_{k,0}|s_1-s_2|}, \tag{16}$$

where $|k\rangle$ is the unbound eigenvector of (11) of eigenvalue $Tk^2/(2\chi)$ and $\Delta\epsilon_{k,0} = Tk^2/(2\chi) - \epsilon_0$, where ϵ_0 is the remaining bound eigenvalue.

From (16), by bringing $\exp(\epsilon_0|\Delta s|)$ out of the integral and noticing that $\exp\left[-T|\Delta s|k^2(2\chi)^{-1}\right] \xrightarrow{|\Delta s|\to\infty} \delta(k)(2\pi\chi\beta/|\Delta s|)^{1/2}$, for large $|\Delta s|$, one finally has

$$G(s_1, s_2) \propto \frac{e^{\epsilon_0|s_1-s_2|}}{\sqrt{s_1 - s_2}}, \tag{17}$$

showing that the correlation length $\xi = -1/\epsilon_0$ approaches infinity exponentially fast at transition:

$$\xi \sim \exp\left(\frac{2\pi}{\sqrt{(\tau/T)^2 - 1/4}}\right). \tag{18}$$

Some similarities are evident in this behavior with the Kosterlitz-Thouless (KT) transition, but this should not be over interpreted, as we are dealing with an externally driven situation. Unlike the KT case, here we do have an order parameter, $\langle\omega\rangle$, which disappears at transition. Also, the external field, τ, breaks the symmetry of the problem. In fact, if we want to establish a formal similarity between the two transitions, we must note that in the formula for the correlation length τ plays the role of T in the KT transition. Similarly, while our heuristic argument in (5) is reminiscent of the analogous one for KT, energy and entropy are exchanged: while in KT the formation of an unbound topological structure, the vortex, is favored by entropy at high T and disfavored by energy, the formation of a helical structure is favored by energy at high τ and disfavored by entropy.

In experiments of single molecule manipulation, one fixes the supercoiling and measures the resulting torque. The transition proposed here can thus be revealed experimentally by reducing the supercoiling to zero and observing that the torque converges very slowly to the value $T/2$, which at room temperature corresponds to ~2 pN×nm. In practice there will also be an inner torsional degree of freedom ζ of stiffness μ. Therefore the total supercoiling will be given by $\sigma = \langle\omega\rangle + \tau/\mu$: experimentally the transition will be manifested by plotting τ/T vs. $\sigma - \tau/\mu$, as shown in Fig. 1.

In conclusion, inspired by a single molecule problem, we have introduced a simple model of a free hamiltonian whose helicity is coupled to a uniform external field, and shown that it leads to an infinite order topological transition. This transition is driven by the competition of entropy and energy, both of which are lowered by helical fluctuations.

Our formalism is more general and might be applied, for instance, to kinetic transitions in problems of Langevin dynamics in an external field $\vec{w}$. Indeed a path integral formalism applies to that case, with lagrangian $(\dot{\vec{x}} - \vec{w})^2/4D - \mathrm{div}\vec{w}/2$ [20]. One can verify that the choice of a vortex $\vec{w} = \tau\hat{e}_3 \wedge \vec{x}/|\vec{x}|^2 + \vec{x}\phi(|\vec{x}|)$, where τ is now the vorticity, can lead to an expression similar to the one in (1) when $\vec{x}$ is represented in polar coordinates, and for appropriate choice of the field ϕ. That would lead to a kinetic transition in τ/D, due to a competition between vorticity and diffusion. Also, $\mathcal{H}$ can be rewritten in terms of a boson condensate as $\beta\mathcal{H} = \chi|\dot{\psi}|^2/2 + i\tau g\left(\psi^*\dot{\psi} - \dot{\psi}^*\psi\right)/2|\psi|^2$. We will discuss these different contents elsewhere.

C. Nisoli is grateful to P. Lammert (Penn State) for discussions. This work was carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DEAC52-06NA25396.

[1] S. F. Edwards, *Proc. Phys. Soc.* **85** 4 (1965)

[2] P. G. De Gennes "Introduction to Polymer Dynamics (Lezioni Lincee)" (Cambridge University Press 1990); "Scaling Concepts in Polymer Physics" (Cornell University Press, 1979)

[3] P. J. Flory "Selected works of Paul J. Flory" (Stanford University Press, 1985).

[4] K. C. Neuman and A. Nagy, *Nature Methods* **5** 491 (2008)

[5] N. C. Seeman, *Nature* **421**, 427 (2003).

[6] S Iijima, *Nature* **354**, 56 (1991). M J. Green, N. Behabtu, M. Pasquali, W. W. Adams, *Polymer* **50** 4979 (2009)

[7] J. F Marko *Europhys. Lett.* **38**, 183 (1997).

[8] S. Cocco and R. Monasson, *Phys. Rev. Lett.* **83**, 5178 (1999).

[9] M. Barbi, S. Cocco, and M. Peyrard, *Phys. Lett. A* **253**, 358 (1999).

[10] C. Nisoli, A. R. Bishop, *Phys. Rev. Lett.* **107**, 068102 (2011).

[11] C. Nisoli *et al.*, *Phys. Rev. Lett.* **104**, 025503 (2010).

[12] T.r. Strick, J. F. Allemand, D. Bensimon, V. Croquette, *Biophys J* **74** 2016 (1998).

[13] C. Bustamante *et al.*, *Nature* **421**, 423 (2003).

[14] Z. Bryant *et al.*, *Nature* **424**, 338 (2003).

[15] K. S. Gupta and S. G. Rajeev, *Phys. Rev. D* **48** 5490 (1993).

[16] H. E. Camblong *et al.*, *Phys. Rev. Lett.* **85**, 1590 (2000).

[17] O. H. Crawford, *Proc. Phys. Soc. London* **91**, 279 (1967), and references therein.

[18] D. Bouazziz, M. Bawin *Phys. Rev. A* **78**, 032110 (2008).

[19] E. Marinari and G. Parisi, *Europhys. Lett.* **15**, 721 (1991).

[20] J. Zinn-Justin, "Quantum field theory and critical phenomena" (Clarendon Press, Oxford, 2002)

[21] Since such experiments can be performed with tensions up to 10^2 pN×nm that suggests a polymer of persistence length of the order of nanometers. But one can consider other special arrangements. Imagine that monomers are connected by a concertina-type hinge, as the one common in articulated drinking straws. Then flexural rigidity will be then zero, whereas inner torsional rigidity can be taken as infinite.

[22] The fact that $\Delta f_{\bar{\omega}}$ becomes unbounded from below in $\bar{\omega}$ when $T > 2|\tau|$ is a consequence of neglecting the cut-off factor g in this heuristic computation.